

Standard order for Adult Sepsis / Septic shock โรงพยาบาลบ้านนา

Progress Note	Date time	Order for one Day	Date time	Order for Continuous										
Early detection -BT > 38 c or < 36 c -HR > 90 /min -RR > 20 or PaCO2 < 32 -WBC > 12000 or < 4000 band form >10% Diagnosis ☐ Sepsis ☐ Septic shock (Dx. At.....น.)		☐ Admit ward..... ☐ H/C x 2 Spec ก่อนให้ATBเวลา..... ☐ Lactate, CBC,BUN,Cr, Electrolyte ☐ LFT ☐ PT/PTT/INR ☐ Ca ,Mg, PO4 (lab emergency) ☐ CXR ☐ EKG 12 lead ☐ DTX stat.....		☐ Diet..... ☐ NPO ☐ Record V/S q.....hr ☐ Record I/O ☐ Retained NG ☐ Retained Foley's catheter ☐ DTX ทุกhr. keep 80- 200 mg%										
Source: ☐ Pneumonia ☐ UTI ☐ Other.....		IV Fluid ☐ RLS ☐ NSS เวลา.....น. ☐ 500 ml. drip in 30 นาที x.....ครั้ง ☐ KVO if volume overload รวม load iv ทั้งหมด.....ml. (Total 30ml/kg in 3 hr.)ในรายที่มี Septic shock ☐ Sputum Gram Stain, C/S ☐ U/A ,U/C ☐ Pus gram stain, culture from..... ☐ อื่นๆ.....		☐ ATB เวลา.....น.(ภายใน 1 ชม.) ☐ Ceftriaxone..... ☐ Ceftazidime..... ☐										
Supportive care : Goal -MAP > 65 mmHg -Urine ≥ 0.5 ml/kg/hr		☐ On ETT no.....ลิ้น.....cm ☐ On oxygen..... Keep O2Sat ≥ 95% Keep MAP > 65 mmHg PR < 140 ☐ Norepinephrine 4 mg+5%DW 250 ml iv rate.....ml/hr. (4 : 250) <table border="1" style="width:100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width:15%;">BW (kg)</td> <td style="width:15%;"><40</td> <td style="width:15%;">40-60</td> <td style="width:15%;">61-80</td> <td style="width:15%;">>80</td> </tr> <tr> <td>Rate(ml/hr)</td> <td>10</td> <td>15</td> <td>20</td> <td>25</td> </tr> </table> titrate ครั้งละ3-5ml/hr.ทุก 15 นาที (max 180 ml/hr) ☐ Dopamine (2 :1) iv rate.....ml/hr. (*กรณี HR < 40) Max30ml/hr.	BW (kg)	<40	40-60	61-80	>80	Rate(ml/hr)	10	15	20	25		☐ Paracetamol (500) 1tab prn. q 4-6 hr. ☐ Omeprazole 40 mg iv OD ☐ If urine output < 120 ml/4 hr. ให้ notify แพทย์
BW (kg)	<40	40-60	61-80	>80										
Rate(ml/hr)	10	15	20	25										
Name of Patient		Age		H.N.										
Department of Service		Ward		Attending Physician										